

Notes on the Dipole Coordinate System

M. Swisdak*

Plasma Physics Division, Naval Research Laboratory, Washington, DC, 20375

Abstract

A strong magnetic field can make it advantageous to work in a coordinate system aligned with dipolar field lines. This monograph collect the formulas for some of the most frequently used expressions and operations in dipole coordinates.

In some physical systems (*e.g.*, the terrestrial ionosphere or the solar corona) the magnetic field can impose a strong anisotropy by restricting transport processes perpendicular to the lines of force. If the field is approximately dipolar it can be useful to work in an aligned coordinate system even though the vector operations are somewhat more complicated than in a Cartesian or spherical polar representation. Here we present several of the formulas that frequently occur when working with vectors in dipolar coordinates. Although some of the results have previously appeared in the literature [2, 4], others seem to be new.

There are many possible (related) choices for the dipolar coordinates. The right-handed orthogonal system considered here, (q, p, ϕ) , is defined in terms of the usual spherical polar coordinates, (r, θ, ϕ) , by

$$q = \frac{\cos \theta}{r^2} \quad p = \frac{r}{\sin^2 \theta} \quad \phi = \phi \quad (1)$$

p is constant along a dipolar field line while q parameterizes the displacement parallel to the field: $q = 0$ at the equator, $q \rightarrow -\infty$ as $\theta \rightarrow \pi$ and $q \rightarrow +\infty$ as $\theta \rightarrow 0$.

1 Inverse Transformation

The inversion of (1) — *i.e.*, finding (r, θ) given (q, p) — involves the solution of a non-trivial equation. Substituting for r gives $qp^2 = \cos \theta / \sin^4 \theta$ while eliminating θ leads to the polynomial expression

$$q^2 r^4 + \frac{r}{p} - 1 = 0 \quad (2)$$

Descartes's rule of signs states that (2) has exactly one positive, real root and, as has been previously noted [3], since (2) is a quartic this root has an algebraic representation. To be useful in numerical models however such a solution has to be expressed in a computationally stable form. Define the auxiliary quantities

$$\alpha = \frac{256}{27} q^2 p^4 \quad \beta = (1 + \sqrt{1 + \alpha})^{2/3} \quad \gamma = \sqrt[3]{\alpha} \quad (3)$$

and

$$\mu = \frac{1}{2} \left(\frac{\beta^2 + \beta\gamma + \gamma^2}{\beta} \right)^{3/2} \quad (4)$$

Then the positive real root of (2) is

$$r = \frac{4\mu}{(1 + \mu)(1 + \sqrt{2\mu - 1})} p \quad (5)$$

*swisdak@ppd.nrl.navy.mil

Since $\mu \geq 1$ this formulation eliminates the possibility of a catastrophic cancellation between terms. To complete the inversion recall that $\sin^2 \theta = r/p$.

Although (5) makes it possible, in principle, to express formulas in terms of either r and θ or q and p , it is usually simpler to use the former representation.

2 Coordinate Derivatives

In this section we ignore the ϕ coordinate. The partial derivatives of the dipolar coordinates with respect to the spherical polar coordinates are

$$\frac{\partial q}{\partial r} = -\frac{2 \cos \theta}{r^3} \quad \frac{\partial p}{\partial r} = \frac{1}{\sin^2 \theta} \quad (6)$$

$$\frac{\partial q}{\partial \theta} = -\frac{\sin \theta}{r^2} \quad \frac{\partial p}{\partial \theta} = -\frac{2r \cos \theta}{\sin^3 \theta} \quad (7)$$

The Jacobian is then

$$\frac{\partial(q, p)}{\partial(r, \theta)} = \frac{\delta^2}{r^2 \sin^3 \theta} \quad (8)$$

where $\delta(\theta) = \sqrt{1 + 3 \cos^2 \theta}$. The derivatives of the spherical polar coordinates with respect to the dipolar coordinates are

$$\frac{\partial r}{\partial q} = -\frac{2r^3 \cos \theta}{\delta^2} \quad \frac{\partial r}{\partial p} = \frac{\sin^4 \theta}{\delta^2} \quad (9)$$

$$\frac{\partial \theta}{\partial q} = -\frac{r^2 \sin \theta}{\delta^2} \quad \frac{\partial \theta}{\partial p} = -\frac{2 \cos \theta \sin^3 \theta}{r \delta^2} \quad (10)$$

A few second derivatives are also occasionally useful

$$\frac{\partial^2 r}{\partial q^2} = -\frac{2r^5}{\delta^6}(1 - 10 \cos^2 \theta - 15 \cos^4 \theta) \quad \frac{\partial^2 r}{\partial p^2} = -\frac{4 \sin^6 \theta \cos^2 \theta}{r \delta^4}(5 + 3 \cos^2 \theta) \quad (11)$$

$$\frac{\partial^2 \theta}{\partial q^2} = \frac{r^4 \sin \theta \cos \theta}{\delta^6}(11 + 9 \cos^2 \theta) \quad \frac{\partial^2 \theta}{\partial p^2} = -\frac{2 \sin^5 \theta \cos \theta}{r^2 \delta^6}(1 - 16 \cos^2 \theta - 9 \cos^4 \theta) \quad (12)$$

3 Unit Vectors

The expressions for the dipolar and spherical polar unit vectors in the alternate coordinate system are

$$\hat{\mathbf{q}} = -\frac{2 \cos \theta}{\delta} \hat{\mathbf{r}} - \frac{\sin \theta}{\delta} \hat{\boldsymbol{\theta}} \quad \hat{\mathbf{p}} = \frac{\sin \theta}{\delta} \hat{\mathbf{r}} - \frac{2 \cos \theta}{\delta} \hat{\boldsymbol{\theta}} \quad (13)$$

$$\hat{\mathbf{r}} = -\frac{2 \cos \theta}{\delta} \hat{\mathbf{q}} + \frac{\sin \theta}{\delta} \hat{\mathbf{p}} \quad \hat{\boldsymbol{\theta}} = -\frac{\sin \theta}{\delta} \hat{\mathbf{q}} - \frac{2 \cos \theta}{\delta} \hat{\mathbf{p}} \quad (14)$$

3.1 First Derivatives of Unit Vectors

$$\frac{\partial \hat{\mathbf{r}}}{\partial q} = \hat{\boldsymbol{\theta}} \frac{\partial \theta}{\partial q} \quad \frac{\partial \hat{\mathbf{r}}}{\partial p} = \hat{\boldsymbol{\theta}} \frac{\partial \theta}{\partial p} \quad \frac{\partial \hat{\mathbf{r}}}{\partial \phi} = \hat{\boldsymbol{\phi}} \sin \theta \quad (15)$$

$$\frac{\partial \hat{\boldsymbol{\theta}}}{\partial q} = -\hat{\mathbf{r}} \frac{\partial \theta}{\partial q} \quad \frac{\partial \hat{\boldsymbol{\theta}}}{\partial p} = -\hat{\mathbf{r}} \frac{\partial \theta}{\partial p} \quad \frac{\partial \hat{\boldsymbol{\theta}}}{\partial \phi} = \hat{\boldsymbol{\phi}} \cos \theta \quad (16)$$

$$\frac{\partial \hat{\boldsymbol{\phi}}}{\partial q} = 0 \quad \frac{\partial \hat{\boldsymbol{\phi}}}{\partial p} = 0 \quad \frac{\partial \hat{\boldsymbol{\phi}}}{\partial \phi} = -\hat{\mathbf{r}} \sin \theta - \hat{\boldsymbol{\theta}} \cos \theta \quad (17)$$

and

$$\frac{\partial \hat{\mathbf{q}}}{\partial q} = \hat{\mathbf{p}} \frac{3(1 + \cos^2 \theta)}{\delta^2} \frac{\partial \theta}{\partial q} \quad \frac{\partial \hat{\mathbf{q}}}{\partial p} = \hat{\mathbf{p}} \frac{3(1 + \cos^2 \theta)}{\delta^2} \frac{\partial \theta}{\partial p} \quad \frac{\partial \hat{\mathbf{q}}}{\partial \phi} = -\hat{\phi} \frac{3 \cos \theta \sin \theta}{\delta} \quad (18)$$

$$\frac{\partial \hat{\mathbf{p}}}{\partial q} = -\hat{\mathbf{q}} \frac{3(1 + \cos^2 \theta)}{\delta^2} \frac{\partial \theta}{\partial q} \quad \frac{\partial \hat{\mathbf{p}}}{\partial p} = -\hat{\mathbf{q}} \frac{3(1 + \cos^2 \theta)}{\delta^2} \frac{\partial \theta}{\partial p} \quad \frac{\partial \hat{\mathbf{p}}}{\partial \phi} = \hat{\phi} \frac{1 - 3 \cos^2 \theta}{\delta} \quad (19)$$

$$\frac{\partial \hat{\phi}}{\partial q} = 0 \quad \frac{\partial \hat{\phi}}{\partial p} = 0 \quad \frac{\partial \hat{\phi}}{\partial \phi} = \hat{\mathbf{q}} \frac{3 \cos \theta \sin \theta}{\delta} - \hat{\mathbf{p}} \frac{1 - 3 \cos^2 \theta}{\delta} \quad (20)$$

3.2 Some Second Derivatives

$$\frac{\partial^2 \hat{\mathbf{q}}}{\partial q^2} = -\hat{\mathbf{q}} \left(\frac{3(1 + \cos^2 \theta)}{\delta^2} \frac{\partial \theta}{\partial q} \right)^2 - \hat{\mathbf{p}} \frac{3r^2 \cos \theta}{\delta^6} (15 + 16 \cos^2 \theta + 9 \cos^4 \theta) \frac{\partial \theta}{\partial q} \quad (21)$$

$$\frac{\partial^2 \hat{\mathbf{q}}}{\partial p^2} = -\hat{\mathbf{q}} \left(\frac{3(1 + \cos^2 \theta)}{\delta^2} \frac{\partial \theta}{\partial p} \right)^2 + \hat{\mathbf{p}} \frac{3 \sin^2 \theta}{r \delta^6} (1 - 23 \cos^2 \theta - 17 \cos^4 \theta - 9 \cos^6 \theta) \frac{\partial \theta}{\partial p} \quad (22)$$

$$\frac{\partial^2 \hat{\mathbf{p}}}{\partial q^2} = -\hat{\mathbf{p}} \left(\frac{3(1 + \cos^2 \theta)}{\delta^2} \frac{\partial \theta}{\partial q} \right)^2 + \hat{\mathbf{q}} \frac{3r^2 \cos \theta}{\delta^6} (15 + 16 \cos^2 \theta + 9 \cos^4 \theta) \frac{\partial \theta}{\partial q} \quad (23)$$

$$\frac{\partial^2 \hat{\mathbf{p}}}{\partial p^2} = -\hat{\mathbf{p}} \left(\frac{3(1 + \cos^2 \theta)}{\delta^2} \frac{\partial \theta}{\partial p} \right)^2 - \hat{\mathbf{q}} \frac{3 \sin^2 \theta}{r \delta^6} (1 - 23 \cos^2 \theta - 17 \cos^4 \theta - 9 \cos^6 \theta) \frac{\partial \theta}{\partial p} \quad (24)$$

$$\frac{\partial^2 \hat{\phi}}{\partial \phi^2} = -\hat{\phi} \quad (25)$$

4 Metric; Differential Line, Area, and Volume Elements

Since the dipole system is orthogonal the only non-zero components of the metric are the diagonal elements (i.e., $g_{ij} = 0$ for $i \neq j$). The associated scale factors (h_i , where $h_i^2 = g_{ii}$) are

$$h_q = \frac{r^3}{\delta} \quad h_p = \frac{\sin^3 \theta}{\delta} \quad h_\phi = r \sin \theta \quad (26)$$

With these we can immediately write down the differential elements

$$d\mathbf{r} = \hat{\mathbf{q}} \frac{r^3}{\delta} dq + \hat{\mathbf{p}} \frac{\sin^3 \theta}{\delta} dp + \hat{\phi} r \sin \theta d\phi \quad (27)$$

$$d\sigma_{qp} = \frac{r^3 \sin^3 \theta}{\delta^2} dq dp \quad d\sigma_{q\phi} = \frac{r^4 \sin \theta}{\delta} dq d\phi \quad d\sigma_{p\phi} = \frac{r \sin^4 \theta}{\delta} dp d\phi \quad (28)$$

$$d\tau = \frac{r^4 \sin^4 \theta}{\delta^2} dq dp d\phi \quad (29)$$

The Christoffel symbols of the second kind, as defined by Arfken [1], are given by the formula

$$\Gamma^m_{ij} = \frac{1}{2} g^{km} \left(\frac{\partial g_{ik}}{\partial x^j} + \frac{\partial g_{jk}}{\partial x^i} - \frac{\partial g_{ij}}{\partial x^k} \right) \quad (30)$$

where $g^{ii} = 1/g_{ii}$. In matrix form they are

$$\Gamma^q = \begin{pmatrix} -\frac{3r^2 \cos \theta}{\delta^4} (3 + 5 \cos^2 \theta) & \frac{3 \sin^4 \theta}{r \delta^4} (1 + \cos^2 \theta) & 0 \\ \frac{3 \sin^4 \theta}{r \delta^4} (1 + \cos^2 \theta) & \frac{6 \sin^6 \theta \cos \theta}{r^4 \delta^4} (1 + \cos^2 \theta) & 0 \\ 0 & 0 & \frac{3 \sin \theta \cos \theta}{r^2} \end{pmatrix} \quad (31)$$

$$\Gamma^p = \begin{pmatrix} -\frac{3r^5}{\delta^4 \sin^2 \theta} (1 + \cos^2 \theta) & -\frac{6r^2 \cos \theta}{\delta^4} (1 + \cos^2 \theta) & 0 \\ -\frac{6r^2 \cos \theta}{\delta^4} (1 + \cos^2 \theta) & -\frac{12 \sin^2 \theta \cos^2 \theta}{r \delta^4} (1 + \cos^2 \theta) & 0 \\ 0 & 0 & \frac{r}{\sin^2 \theta} (1 + \cos^2 \theta) \end{pmatrix} \quad (32)$$

$$\Gamma^\phi = \begin{pmatrix} 0 & 0 & -\frac{3r^2 \cos \theta}{\delta^2} \\ 0 & 0 & \frac{\sin^2 \theta}{r \delta^2} (1 - 3 \cos^2 \theta) \\ -\frac{3r^2 \cos \theta}{\delta^2} & \frac{\sin^2 \theta}{r \delta^2} (1 - 3 \cos^2 \theta) & 0 \end{pmatrix} \quad (33)$$

5 Vector Operations

The differential operators can be derived from the metric tensor. In what follows f is a scalar, $\mathbf{A}$ and $\mathbf{B}$ are vectors, and $\mathbf{T}$ is a tensor.

5.1 Gradient

$$\nabla f = \hat{\mathbf{q}} \frac{\delta}{r^3} \frac{\partial f}{\partial q} + \hat{\mathbf{p}} \frac{\delta}{\sin^3 \theta} \frac{\partial f}{\partial p} + \hat{\phi} \frac{1}{r \sin \theta} \frac{\partial f}{\partial \phi} \quad (34)$$

5.2 Divergence

$$\nabla \cdot \mathbf{A} = \frac{\delta^2}{r^6} \frac{\partial}{\partial q} \left(\frac{r^3}{\delta} A_q \right) + \frac{\delta^2}{r^4 \sin^4 \theta} \frac{\partial}{\partial p} \left(\frac{r^4 \sin \theta}{\delta} A_p \right) + \frac{1}{r \sin \theta} \frac{\partial A_\phi}{\partial \phi} \quad (35)$$

which can also be written as either

$$\nabla \cdot \mathbf{A} = \frac{\delta^2}{r^6} \frac{\partial}{\partial q} \left(\frac{r^3}{\delta} A_q \right) + \frac{\delta^2}{\sin^6 \theta} \frac{\partial}{\partial p} \left(\frac{\sin^3 \theta}{\delta} A_p \right) + \frac{4}{r \delta \sin \theta} A_p + \frac{1}{r \sin \theta} \frac{\partial A_\phi}{\partial \phi} \quad (36)$$

or

$$\nabla \cdot \mathbf{A} = \frac{\delta}{r^3} \frac{\partial A_q}{\partial q} - \frac{3 \cos \theta}{r \delta^3} (3 + 5 \cos^2 \theta) A_q + \frac{\delta}{\sin^3 \theta} \frac{\partial A_p}{\partial p} + \frac{4}{r \delta^3 \sin \theta} (1 - 3 \cos^4 \theta) A_p + \frac{1}{r \sin \theta} \frac{\partial A_\phi}{\partial \phi} \quad (37)$$

5.3 Curl

$$\begin{aligned} \nabla \times \mathbf{A} = & \hat{\mathbf{q}} \frac{1}{r \sin \theta} \left[\frac{\delta}{\sin^3 \theta} \frac{\partial}{\partial p} (r \sin \theta A_\phi) - \frac{\partial A_p}{\partial \phi} \right] \\ & + \hat{\mathbf{p}} \frac{1}{r \sin \theta} \left[\frac{\partial A_q}{\partial \phi} - \frac{\delta}{r^3} \frac{\partial}{\partial q} (r \sin \theta A_\phi) \right] \\ & + \hat{\phi} \frac{\delta^2}{r^3 \sin^3 \theta} \left[\frac{\partial}{\partial q} \left(\frac{\sin^3 \theta}{\delta} A_p \right) - \frac{\partial}{\partial p} \left(\frac{r^3}{\delta} A_q \right) \right] \end{aligned} \quad (38)$$

which is equivalent to

$$\begin{aligned} \nabla \times \mathbf{A} = & \hat{\mathbf{q}} \left[\frac{\delta}{\sin^3 \theta} \frac{\partial A_\phi}{\partial p} + \frac{1 - 3 \cos^2 \theta}{r \delta \sin \theta} A_\phi - \frac{1}{r \sin \theta} \frac{\partial A_p}{\partial \phi} \right] \\ & + \hat{\mathbf{p}} \left[\frac{1}{r \sin \theta} \frac{\partial A_q}{\partial \phi} - \frac{\delta}{r^3} \frac{\partial A_\phi}{\partial q} + \frac{3 \cos \theta}{r \delta} A_\phi \right] \\ & + \hat{\phi} \left[\frac{\delta}{r^3} \frac{\partial A_p}{\partial q} - \frac{6 \cos \theta}{r \delta^3} (1 + \cos^2 \theta) A_p - \frac{\delta}{\sin^3 \theta} \frac{\partial A_q}{\partial p} - \frac{3 \sin \theta}{r \delta^3} (1 + \cos^2 \theta) A_q \right] \end{aligned} \quad (39)$$

5.4 Scalar Laplacian

$$\nabla^2 f = \frac{\delta^2}{r^6} \frac{\partial^2 f}{\partial q^2} + \frac{\delta^2}{r^4 \sin^4 \theta} \frac{\partial}{\partial p} \left(\frac{r^4}{\sin^2 \theta} \frac{\partial f}{\partial p} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 f}{\partial \phi^2} \quad (40)$$

which can also be written as

$$\nabla^2 f = \frac{\delta^2}{r^6} \frac{\partial^2 f}{\partial q^2} + \frac{\delta^2}{\sin^6 \theta} \frac{\partial^2 f}{\partial p^2} + \frac{4}{r \sin^4 \theta} \frac{\partial f}{\partial p} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 f}{\partial \phi^2} \quad (41)$$

5.5 Vector Laplacian

$$\begin{aligned} \nabla^2 \mathbf{A} = & \hat{\mathbf{q}} \left[\nabla^2 A_q + \frac{6 \sin \theta}{r^4 \delta^2} (1 + \cos^2 \theta) \frac{\partial A_p}{\partial q} + \frac{12 \cos \theta}{r \delta^2 \sin^3 \theta} (1 + \cos^2 \theta) \frac{\partial A_p}{\partial p} + \frac{6 \cos \theta}{r^2 \delta \sin \theta} \frac{\partial A_\phi}{\partial \phi} \right. \\ & \left. - \frac{9}{r^2 \delta^4} (1 + 3 \cos^2 \theta + 4 \cos^4 \theta) A_q - \frac{12 \cos \theta}{r^2 \delta^4 \sin \theta} (1 + 3 \cos^4 \theta) A_p \right] \\ & + \hat{\mathbf{p}} \left[\nabla^2 A_p - \frac{6 \sin \theta}{r^4 \delta^2} (1 + \cos^2 \theta) \frac{\partial A_q}{\partial q} - \frac{12 \cos \theta}{r \delta^2 \sin^3 \theta} (1 + \cos^2 \theta) \frac{\partial A_q}{\partial p} - \frac{2}{r^2 \delta \sin^2 \theta} (1 - 3 \cos^2 \theta) \frac{\partial A_\phi}{\partial \phi} \right. \\ & \left. + \frac{18 \sin \theta \cos \theta}{r^2 \delta^4} (1 + \cos^2 \theta) A_q - \frac{2}{r^2 \delta^4 \sin^2 \theta} (5 + 3 \cos^2 \theta - 9 \sin^2 \theta \cos^4 \theta) A_p \right] \\ & + \hat{\phi} \left[\nabla^2 A_\phi - \frac{6 \cos \theta}{r^2 \delta \sin \theta} \frac{\partial A_q}{\partial \phi} + \frac{2}{r^2 \delta \sin^2 \theta} (1 - 3 \cos^2 \theta) \frac{\partial A_p}{\partial \phi} - \frac{A_\phi}{r^2 \sin^2 \theta} \right] \end{aligned} \quad (42)$$

5.6 Directional Derivative

$$\begin{aligned} (\mathbf{A} \cdot \nabla) \mathbf{B} = & \hat{\mathbf{q}} \left[\mathbf{A} \cdot \nabla B_q + \frac{3 \sin \theta}{r \delta^3} (1 + \cos^2 \theta) A_q B_p + \frac{6 \cos \theta}{r \delta^3} (1 + \cos^2 \theta) A_p B_p + \frac{3 \cos \theta}{r \delta} A_\phi B_\phi \right] \\ & + \hat{\mathbf{p}} \left[\mathbf{A} \cdot \nabla B_p - \frac{3 \sin \theta}{r \delta^3} (1 + \cos^2 \theta) A_q B_q - \frac{6 \cos \theta}{r \delta^3} (1 + \cos^2 \theta) A_p B_q \right. \\ & \left. - \frac{1}{r \delta \sin \theta} (1 - 3 \cos^2 \theta) A_\phi B_\phi \right] \\ & + \hat{\phi} \left[\mathbf{A} \cdot \nabla B_\phi - \frac{3 \cos \theta}{r \delta} A_\phi B_q + \frac{1}{r \delta \sin \theta} (1 - 3 \cos^2 \theta) A_\phi B_p \right] \end{aligned} \quad (43)$$

5.7 Divergence of a Tensor

$$\begin{aligned} \nabla \cdot \mathbf{T} = & \hat{\mathbf{q}} \left[\nabla \cdot (\mathbf{T}_{qq} \hat{\mathbf{q}} + \mathbf{T}_{pq} \hat{\mathbf{p}} + \mathbf{T}_{\phi q} \hat{\phi}) + \frac{3 \sin \theta}{r \delta^3} (1 + \cos^2 \theta) \mathbf{T}_{qp} + \frac{6 \cos \theta}{r \delta^3} (1 + \cos^2 \theta) \mathbf{T}_{pp} + \frac{3 \cos \theta}{r \delta} \mathbf{T}_{\phi \phi} \right] \\ & + \hat{\mathbf{p}} \left[\nabla \cdot (\mathbf{T}_{qp} \hat{\mathbf{q}} + \mathbf{T}_{pp} \hat{\mathbf{p}} + \mathbf{T}_{\phi p} \hat{\phi}) - \frac{3 \sin \theta}{r \delta^3} (1 + \cos^2 \theta) \mathbf{T}_{qq} - \frac{6 \cos \theta}{r \delta^3} (1 + \cos^2 \theta) \mathbf{T}_{pq} \right. \\ & \left. - \frac{1}{r \delta \sin \theta} (1 - 3 \cos^2 \theta) \mathbf{T}_{\phi \phi} \right] \\ & + \hat{\phi} \left[\nabla \cdot (\mathbf{T}_{q\phi} \hat{\mathbf{q}} + \mathbf{T}_{p\phi} \hat{\mathbf{p}} + \mathbf{T}_{\phi\phi} \hat{\phi}) - \frac{3 \cos \theta}{r \delta} \mathbf{T}_{\phi q} + \frac{1}{r \delta \sin \theta} (1 - 3 \cos^2 \theta) \mathbf{T}_{\phi p} \right] \end{aligned} \quad (44)$$

Acknowledgments

This work was supported by the Office of Naval Research.

References

- [1] Arfken, G., *Mathematical Method for Physicists, 3rd ed.*, San Diego, CA: Academic Press, pp. 158-163, 1985.
- [2] Fatkullin, M. N., and Yu. S. Sitnov, Dipole coordinate system and some of its characteristics, *Geomagnetism and Aeronomy*, 12, 293–295, 1972, translated from Russian.
- [3] Kageyama, A., T. Sugiyama, K. Watanabe, and T. Sato, A note on the dipole coordinates, *Computers and Geosciences*, 32, 265–269, 2006.
- [4] Orens, J. H., T. R. Young, Jr., E. S. Oran, and T. P. Coffey, Vector operations in a dipole coordinate system, *NRL Memo. Rep. 3984*, Naval Research Laboratory, Washington, D.C., 1979.